

Report on the first phytoPal meeting

University of Leicester, 11-12th December 2003



Delegates (from left to right). Alex Page, Catherine Duggan, Chris Peat, Gary Mullins, Thomas Servais, Dick Aldridge, Malgorzata Moczydlowska-Vidal, Monika Masiak, Alain Le Hérissé, Reed Wicander, Elena Raevskaya, Marco Vecoli, Li Jun, Ian Troth, John Marshall and Ken Dorning (photographer).

In 2003 the Leverhulme Trust awarded a Research Interchange grant to Dick Aldridge (University of Leicester). The objective of this grant is to foster closer links between scientists interested in Palaeozoic palynology and principally the study of the acritarchs and prasinophycean algae. The name phytoPal was, therefore, coined. The free exchange of ideas lies at the heart of phytoPal. This exchange takes place via regular workshop meetings, through discussion on an email distribution list and through exchange visits.

Thursday 11th December Management Meeting

Attended by Dick Aldridge, Ken Dorning, Alain Le Hérissé, Li Jun, Malgorzata Moczydlowska-Vidal, Gary Mullins, Thomas Servais, Reed Wicander.

Reporting

The Leverhulme Trust require an annual report on both the scientific and financial activity and a list of publications arising out of the project. A final report of 2000 words is required which details the history, activities, dissemination of results and references.

The phytoPal Database

It was agreed that a phytoPal database should be constructed and that this should be a relational database, probably in Microsoft Access 2000. Points raised during discussion on the nature and purpose of the database are outlined below:

- The taxonomy part of the database would be comprehensive and will include all acritarch and prasinophyte taxa even if they were restricted to the Mesozoic or Cenozoic.
- A major aim discussed was a Sepkoski-type curve which can be compared with the distribution of macrofossils etc. It was agreed that this curve would be published as a joint paper. However, it was noted that these types of curve can be related to palaeoenvironmental changes. In addition it was noted that people could use the database to provide a context in which to examine particular parts of the Sepkoski-type curve on an individual basis.
- The database would include detailed information from the Late Neoproterozoic Ediacaran Period (from c. 570 Ma) to the start of the Triassic. This avoids problems with the correlation of the Precambrian-Cambrian boundary. It was decided to begin this aspect of the project at the base of the Cambrian and work back into the Neoproterozoic. Further, it was noted that by including the Neoproterozoic we would encourage links between Precambrian and Palaeozoic phytoplankton workers.
- All phytoplankton and associated groups would be included in the database. These include acritarchs, prasinophytes, zygnemataleans, mazuelloids and calcispheres. In addition multicellular phytoplankton (e.g. *Botryococcus*) and picoplankton would be included where possible. Problems with picoplankton recovery were noted, but it was agreed that this was a topic with immediate research potential.
- Problems were also noted with leiosphaerid taxonomy. The database would allow analyses that included and omitted leiospheres; leiosphere distribution could also be examined independently to test for taxonomic skewing.
- It was decided to retain ecophenotypes as independent morphotaxa.
- The original species identification will be retained in the database even if that identification has been questioned.
- It was noted that the phytoPal database will provide an indirect record of phytoplankton change related to productivity, sedimentation rate, palaeoenvironment and life cycle.
- It might be revealing to examine the distribution of morphologically conservative groups (e.g. *Veryhachium*, *Micrhystridium*) through time.
- The construction of the database would be led by the specialists
- Separate categories will be created for disregarded occurrences of taxa (these decisions will be based on specialist knowledge).
- Possibilities of observing phytoplankton changes related to the shifting positions of palaeogeographical areas through different climatic belts were noted.
- The stratigraphical resolution would be decided on a system-by-system basis depending on how well established the subdivisions were.
- Only the best quality data would be added to the database.
- A cladistic analysis is a possibility and it might be possible to identify those morphological criteria which are most suitable for taxonomic subdivision. At

present there are c. 500 morphological terms in circulation, although they are not consistently used.

Future Workshops

A 'mini-workshop' would be held at the IPC meeting in Granada (July 4th-9th). If possible, funds would be made available for the hire of a venue and a dinner.

The 2004 workshop would be held in conjunction with the Palaeontological Association meeting in Lille and would be hosted by Thomas Servais.

The possibility of holding the 2005 workshop in conjunction with the AASP St. Louis meeting was raised.

A final (unfunded) workshop would be held in conjunction with the 2006 CIMP General Meeting in Prague, Czech Republic.

Publications

The possibility of producing a single large comprehensive publication at the end of the project was discussed and supported. Smaller publications on regional topics would be produced throughout the term of the project.

phytoPal Workshop Leicester 2003

Programme

Friday 12th December The phytoPal Workshop

In the Sparkenhoe & Goscote Room, 4th floor Charles Wilson Building, University of Leicester

- 09:30 **Dick Aldridge**. *University of Leicester*
Welcome
- 09:40 **Thomas Servais**. *Université des Sciences et Technologies de Lille*
Ordovician phytoplankton (acritarch) diversity
- 10:05 **Marco Vecoli**. *Université de Bretagne Occidentale and CNRS*
Biodiversity patterns of Ordovician marine microphytoplankton: biological and palaeoecological meaning
- 10:30 **Reed Wicander**. *Central Michigan University*
Acritarch and prasinophyte diversity trends during the Devonian
- 10:55 tea + coffee
- 11:15 **Catherine Duggan**. *Trinity College Dublin*
Determination of organic maturity based on quantitative determination of acritarch colour
- 11:40 **Ken Dorning**. *University of Sheffield and Pallab Research*
The palynological record: variability in palynomorph preservation, geothermal alteration, sampling strategies, processing methods, recording, techniques, data analysis and palaeoenvironmental interpretation
- 12:05 **Gary Mullins**. *University of Leicester*
The phytopal database
- 12:30 lunch
- 13:30 **Elena Raevskaya**^{1,2} & **Elena Golubkova**¹. 1, *All-Russian Petroleum Research Exploration Institute (VNIGRI)*. 2, *Université des Sciences et Technologies de Lille*
Middle-Upper Cambrian acritarch community from Arctic Russia
- 13:55 **Li Jun**. *Nanjing Institute of Geology and Palaeontology*
Acritarch research in China
- 14:20 **Monika Masiak**¹, **Teresa Podhalańska**² **Marzena Stempień-Sałek**¹. 1, *Polish Academy of Sciences, Institute of Geological Sciences*. 2, *Polish Geological Institute*
Integrated biostratigraphy (acritarch and graptolite) on Ordovician - Silurian boundary in the Bardo Syncline, Holy Cross Mountains, Poland
- 14:45 **Ken Dorning**. *University of Sheffield and Pallab Research*
High resolution palynological data – implications for the interpretation of palaeoenvironmental and climatic changes in the sedimentary record
- 15:10 tea+coffee
- 15:30 **Alain Le Hérisse**. *Université de Bretagne Occidentale*
Evolution of biodiversity in the marine phytoplankton: from acritarchs to dinocysts
- 15:55 **Ian Troth**. *Southampton Oceanography Centre*
Sea levels, paleoclimate and hydrocarbon systems in Bolivia
- 16:20 **Malgorzata Moczydlowska**¹, **Sebastian Willman**¹, **Yin Leiming**². 1, *Uppsala University*. 2, *Nanjing Institute of Geology and Palaeontology*
Life cycle of Early Ordovician acritarch species
- 16:45 **T. Servais**¹, **D.A.T. Harper**², **J. Li**³, **A. Munnecke**⁴, **A.W. Owen**⁵ & **P.M Sheehan**⁶. 1, *Université des Sciences et Technologies de Lille*. 2, *Geologisk Museum, København*. 3, *Institute of Geology and Palaeontology, Nanjing*. 4, *Institut für Paläontologie, Universität Erlangen*. 5, *Division of Earth Sciences, Centre for Geosciences, University of Glasgow*. 6, *Geology Department, Milwaukee Public Museum*.
Proposal of a new IGCP project: Ordovician Palaeogeography and Palaeoclimate: IGCP project n° 503

phytoPal Workshop Leicester 2003

Abstracts

The palynological record: variability in palynomorph preservation, geothermal alteration, sampling strategies, processing methods, recording techniques, data analysis and palaeoenvironmental interpretation

Ken J. Dorning

Pallab Research, 58 Robertson Road, Sheffield S6 5DX, England and Palynology Research – Department of Geography, University of Sheffield. K.J.Dorning@sheffield.ac.uk

The preserved palynomorphs in sedimentary sequences are influenced by depositional processes and early sedimentary diagenesis. With later deep burial, geothermal alteration may also degrade the recoverable palynomorphs. Sampling strategies often sample a small proportion of the available material, with the aim of providing a representative suite of samples. While the basic palynological processing techniques regularly used are broadly similar, the slight differences in method can result in surprisingly different results. Recording techniques are somewhat variable, depending in part on the required data analysis and palaeoenvironmental interpretation. All of these variables will have an influence over the accuracy of palynomorph data records throughout the Palaeozoic.

Talk

High resolution palynological data – implications for the interpretation of palaeoenvironmental and climatic changes in the sedimentary record

Ken J. Dorning

Pallab Research, 58 Robertson Road, Sheffield S6 5DX, England and Palynology Research – Department of Geography, University of Sheffield. K.J.Dorning@sheffield.ac.uk

The absolute abundance of acritarchs in the Palaeozoic varies with sedimentation rate related to sediment type, as well as nutrient availability, which is influenced by palaeoenvironment and palaeolatitude. Analysis of the absolute abundance of the marine acritarch phytoplankton record in Silurian tropical open marine shelf sequences with apparently continuous slow sedimentation show a remarkable consistency in the preserved acritarch assemblages over short distances laterally at the same horizon, with greater differences between stratigraphical samples. The high resolution data may be used to estimate the reliability of low resolution presence / apparent absence data when used in palaeoenvironmental interpretation.

Talk

Acritarch evidence of marine Silurian cyclicity and climate change

Ken J. Dorning¹ & Craig Harvey^{1, 2}

1, Pallab Research, 58 Robertson Road, Sheffield S6 5DX, England and Palynology Research – Department of Geography, University of Sheffield. 2, Shell Expro, Stavanger. K.J.Dorning@sheffield.ac.uk

Silurian marine sedimentary sequences in eastern Avalonia and Baltica show cyclic variation in carbonate content, that can be assigned to more argillaceous Primo episodes, and more calcareous Secundo episodes. In the type Wenlock area, acritarch assemblages of Primo episodes have diverse acritarch assemblages, often with a high relative abundance of *Diexallophasis*. In contrast, Secundo episodes have acritarch assemblages including the very large acritarchs *Estiastra*, *Hoegklintia* and *Pulvinosphaeridium*. Abundances range between 200 and 2000 per gram, with the four most common genera represented by *Leiosphaeridia* at 6-67%, *Veryhachium* at 10-28%, *Micrhystridium* at 7-53% and *Diexallophasis* at 3-37%. Given an averaged sedimentary accumulation rate of 21.5 years per mm, the preserved marine phytoplankton productivity ranges between 25,000 and 250,000 for each square metre per year.

Poster

Determination of organic maturity based on quantitative determination of acritarch colour

Catherine Duggan

Department of Geology, Trinity College, Dublin. duggancm@tcd.ie

The objective of this project is to develop an inexpensive alternative method for the assessment of organic maturity of vitrinite-deficient marine hydrocarbon source rocks, based on colour change in acritarchs. Previously, visual estimation of palynomorph colour has been used to assess maturity, but quantitative colour analysis of palynomorph colour is a more precise and objective method of determining this characteristic. Colour is relatively simple to measure using readily available equipment, the price of which had been steadily falling.

Vitrinite reflectance is the standard technique for assessing the organic maturity of sedimentary rocks. However, very little vitrinite is found in pre-Devonian successions and marine hydrocarbon source rocks of Devonian and younger age commonly contain little vitrinite. In contrast, these lithologies often contain acritarchs and these change colour with increasing temperature. Quantitative determination of the colour of these palynomorphs, calibrated against vitrinite reflectance offers an improved method of assessing sample maturity.

Calibration of acritarch colour against vitrinite reflectance data is currently taking place using Late Devonian shale samples collected from the Midwest USA. These samples contain abundant acritarchs and vitrinite, and range from immature to marginally post-mature. Mean random vitrinite reflectance (R_r) and acritarch colour (relative RGB intensity) determinations are being made in TCD using existing digital photomicrographic and image analysis equipment on fossil plant material already

extracted from the samples. Methods of enabling inter-laboratory calibration are also being explored.

Talk

Evolution of diversity in the marine phytoplankton: from acritarchs to dinocysts

Alain Le Hérisse

UMR 6538 du CNRS, Université de Bretagne Occidentale, 6 Avenue V. Le Gorgeu, BP 809, 29285 BREST Cedex, France. alain.le.herisse@univ-brest.fr

Major changes in the hierarchy of dominant components of the marine microphytoplankton correspond to major events in the biological history of oceans. It is reflected for example by the apparent taxonomic replacement between acritarchs well known from the Proterozoic to Paleozoic and dinocysts from the Mesozoic. Series of observations/new questions are proposed herein to evaluate the prehistory of dinoflagellates and possible links with producers of acritarchs. The possibilities of use of acritarchs are equivalent to those of dinoflagellate workers : to investigate the extent to which the composition of the associations and their distribution may reflect hydrographic characteristics and how it is possible to develop the study of the cysts as paleoceanographic indicators. The best information provided by diversity is obtain when we concentrate the analysis on the level of species and ecosystems. And here, the reappraisal of evolution of diversity in the Phanerozoic microphytoplankton is better understand through precise examples : operational palynology in the Silurian of Saudi Arabia ; palynological tracers of glaciation and deglaciation in the late Ordovician ; phytoplankton development in the late Devonian as a possible key for the modern associations.

Talk

Acritarch research in China

Li Jun

Institute of Geology and Palaontology, Chinese Academy of Sciences (NIGPAS), Chi-Ming-Ssu, Nanjing 210008, China. li_jun54@hotmail.com

Since the first paper published in 1974, more than 100 scientific articles dealing with Chinese Paleozoic acritarchs have been published. These papers have mostly been submitted to Chinese journals and are generally published in Chinese. Most articles deal with the Ordovician acritarchs from Southern China. These results are becoming increasingly important, not only for a better understanding of acritarch biostratigraphy at a local scale, but also for international corration, for global paleobiogeographical scenarios and for paleoecological interpretation. A number of palaeobiogeographical models for Ordovician organic-walled microphytoplankton (acritarchs, prasinophytes, and related groups) have been published during the past 30 years. In general, acritarch workers have recognized two distinct geographic acritarch assemblages in the Ordovician that were attributed to cold and warm-water environments. We present a series of new plots on recently compiled palaeogeographical maps. The review of the literature and the new plots indicate that a number of preliminary conclusions can be drawn. Following minor biogeographical differentiation of acritarch assemblages during the Cambrian, 'provincialism' started at the Cambrian-

Ordovician boundary. In the late Tremadocian a warm-water assemblage seems to be limited to low latitudes localities. From the late Tremadocian and throughout most of the Arenigian an acritarch assemblage following the southern margin of the supercontinent Gondwana can be recognized. It seems reasonable to consider the acritarchs of the palaeocontinent Baltica, at intermediate latitudes, as belonging to a temperate water 'province', which was probably not restricted to the palaeocontinent of Baltica but had a wider distribution at about the same latitude, as some of the elements recorded from Baltica also occur in South China and Argentina. Palynological investigations from several sections located in different parts of the platform allow the observation of an inshore-offshore trend in the late Early Ordovician (Yushmanian) *Didymograptus deflexus* graptolite Biozone and in the early Middle Ordovician (Dawanian) *Azygograptus suecicus* graptolite Biozone. From west to east the acritarch assemblages show not only a remarkable increase in both abundance and diversity with the deepening of the water column, but also changes in the composition of the assemblages. Poorly diversified in the western part of the platform (Kunming area), the acritarchs are more abundant and diversified in the central (Tongzi area) and eastern part (Yichang area) of the Platform. These differences very probably reflect an inshore-offshore trend, and not only palaeogeographical changes or sea-level fluctuations, as previously suggested in the literature.

Talk

Life cycle of Early Ordovician acritarch species

Malgorzata Moczydlowska-Vidal¹, Sebastian Willman¹, Yin Leiming²

1, Uppsala University, Earth Sciences, Palaeobiology, Norbyvägen 13, 756 54 Uppsala, Sweden. 2, Nanjing Institute of Geology and Palaeontology, Academia Sinica, 39 East Beijing Road, Nanjing 21008, China. malgo.vidal@pal.uu.se

Acritarchs are marine planktonic autotrophic protists of heterogeneous origins. Their diversification into many morphotypes occurred throughout Neoproterozoic and Early Palaeozoic; some morphotypes represent green algal classes but most are of unknown biological affinities. Palaeobiology and relationships to extant microbiota of some acritarchs, with emphasis on their life cycle, reproduction and environmental adaptations, may be inferred from phenetic morphological features and cell wall ultrastructure. Microfossils from the Cambrian-Ordovician of China are studied to reveal the wall ultrastructure of vegetative cells and dormant/reproductive cysts, the structural complexity of early eukaryotic cytoskeleton, and to recognize by morphological and ultrastructural means the relationships between various phenotypes. Acritarchs are considered to be preservable cysts of unicellular algae. The new discovery of the entire organism consisting of vegetative envelope and internal cyst shows that some taxa indeed represent the dormant/reproductive cysts whereas other may represent vegetative cells in their complex life cycle. Formation of the cyst, the excystment structure (pylome) and change of the generations (sexual and asexual) in the life cycle of unicellular microbiota may shed light on the development of the early adaptations to survive ecological crises events and as a competitive advantage in increasingly complex marine ecosystems.

Talk

Integrated biostratigraphy (acritarch and graptolite) on Ordovician–Silurian boundary in the Bardo Syncline, Holy Cross Mountains, Poland

Monika Masiak¹, Teresa Podhalańska² Marzena Stempień-Sałek¹

1, Polish Academy of Sciences, Institute of Geological Sciences, Twarda 51/55, 00-818 Warszawa, mmasiak@twarda.pan.pl; mstempie@twarda.pan.pl. 2, Polish Geological Institute, Rakowiecka 4, 00-975 Warszawa, tpod@pgi.waw.pl.

In the southern part of Holy Cross Mountains, in the Bardo Syncline two sections were investigated near the Ordovician–Silurian (O/S) boundary. A conformable and gradual transition from sandy through silty to clayey strata as well as a gradual change of colour of the rock from light to dark occurs in one of them (Bardo Stawy section). That succession indicates a gradual stagnation and progressive oxygen deficiency of the depositional environment related to a deepening of the sea during the post-glacial sea-level rise.

In the second section (Zalesie Nowe) the pass from Ordovician to Silurian occurs within the interval of clayey shales.

Three graptolite biozones are distinguished for the O/S boundary interval deposits: *?persculptus*, *ascensus-acuminatus* and *vesiculosus*. The first, rare graptolites appear below “graptolitic shales” in the *?persculptus* Biozone. The diversity of the assemblage increases in the next two biozones.

An analysis of acritarch frequency shows a decrease of acritarch frequency at the O/S boundary and then a gradual increase in the lower part of the *ascensus-acuminatus* Biozone with a maximum in the *vesiculosus* Biozone. The peak in acritarch frequency noted in the lower part of the *ascensus-acuminatus* Biozone coincides with a considerable increase in Chitinozoa abundance whereas bioturbation was observed slightly below.

The peak in acritarch frequency noted in the lower part of the *vesiculosus* Biozone coincides with a huge increase in Chitinozoa abundance, and the occurrence of bioturbation, brachiopods, and conodonts.

The taxonomic diversity of the acritarch assemblage shows a similar trend although the maximum of acritarch frequency observed in the *vesiculosus* Biozone do not coincide the maximum of acritarch taxonomic diversity.

The change in acritarch assemblages is not related to lithology. The distinct drop in frequency in the *?persculptus* and lower part of *ascensus-acuminatus* biozones occurs in the Zalesie Nowe section within the interval of clayey shales, and, in the Bardo Stawy section, it occurs in a gradual change in lithology (from a sandy and silty to a clayey succession).

Talk

The phytoPal database

Gary Mullins

Department of Geology, University of Leicester, University Road, Leicester LE1 7RH.
glm2@le.ac.uk

The phytoPal (relational) database is currently being constructed within Microsoft Access 2000. The database will contain details of all of the published phytoplankton genera and species. At present the details (name, author, stratigraphical range and taxonomic information) of 724 genera and approximately 1500 species have been entered. However, to fully understand the distribution of the phytoplankton it will be necessary to include additional information on the collector/researcher, geographical locality, palaeoenvironment, lithology, thermal maturation and processing method amongst others. Each of these is likely to introduce its own bias to the data and each will need to be quantified.

A relational database, such as Access, allows data to be 'mined' effectively. For example, queries could be run that would identify those taxa which are restricted to certain stratigraphical intervals, palaeogeographical areas, or palaeoenvironments. Relational databases also provide a good means of making the data public via the WWW. Given that Fensome et al. is now 13 years out of date, part of the database could be made freely available to provide a valuable reference of up to date taxonomic information.

Talk

Acritarch biostratigraphy in the Cambrian-Ordovician of Arctic Russia

Elena Raevskaya^{1,2} & Elena Golubkova¹

1, All-Russian Petroleum Research Exploration Institute (VNIGRI), Litejny pr., 39, St.-Petersburg 191104, Russia. 2, UPRESA 8014 du CNRS (Laboratoire de Paléontologie et Paléogéographie du Paléozoïque), Université des Sciences et Technologies de Lille, Villeneuve d'Ascq Cedex, F-59655, France. elena_raevskaya@hotmail.com

The most comprehensive study on Cambrian and Ordovician acritarchs in Russia had been done in Moscow sineclise and its adjacent areas to the northwest. Only few data regarding the Cambrian–Ordovician are known from the other parts of the large territory. In frame of the Intas–Nemlor (Northern Eurasian Margin and Lomonosov Ridge) project concentrated on magmatism, palaeomagnetism, structural geology and stratigraphy of the Eurasian high Arctic the first palynological study of Lower Palaeozoic strata had been carried out in the central part of Severnaya Zemlya Archipelago (October Revolution Island). Two Cambrian (Universitetskaya and Kurchavinskaya) and two Ordovician (Kruzhikikhskaya and Ushakovskaya) Formations, consisting mostly of terrigenous rocks, were sampled along Kurchavaya and Kanyon rivers. These sediments are exposed in individual blocks that cause difficulty to consecutive investigation. More than 40 samples have been treated to clarify the age of the formations. Among them less than half yielded more or less diverse acritarchs of moderate or poor preservation. However, most of the identified taxa showed high biostratigraphical utility. Five acritarch assemblages had been distinguished in the studied sequences. The oldest one including *Eliasum Ilaniscum*, *Comasphaeridium* sp., *Timofeevia* sp. A, *Timofeevia janischewskyi*, *Cristallinium* sp.,

Retisphaeridium dichamerum, *R. howellii*, suggests a correlation of the rocks with the *Paradoxides paradoxissimus* – *P. forchhammeri* trilobite Zones of Middle Cambrian age. Assemblage II, comprising *Leiofusa stoumonensis*, *Vulcanisphaera turbata*, *C. randomense*, *Cristallinium* cf. *cambriense*, *Timofeevia lancarae* and *T. pentagonalis*, undoubtedly indicates on Upper Cambrian age, being an equivalent to the *Olenus-Parabolina spinulosa* trilobite Zones. The two younger assemblages III and IV are different only in its diversity (that is probably due to the different facial conditions), both distinguished by the appearance of diacrodian (*Acanthodiacrodium*, *Actinothodissus*, *Dasydiacrodium*) and galeate (*Cymatiogalea*, *Stelliferidium*) acritarchs together with *Lusatia dramatica*, *L. triangularis*, *Impluviculus villosiusculus*. This allows a correlation of the sediments with the upper part of Upper Cambrian embracing an interval of the *Protopeltura praecursor*, *Peltura minor* and *P. scarabaeoides* trilobite Zones. None of the Cambrian taxa extends into the youngest assemblage V. The latter is remarkably poor and contains no biostratigraphical markers. The studied succession straddling the Cambrian–Ordovician boundary shows a break in the Late Cambrian stratigraphy of Severnaya Zemlya.

Talk

Ordovician phytoplankton (acritarch) diversity

Thomas Servais

UPRESA 8014 du CNRS (Laboratoire de Paléontologie et Paléogéographie du Paléozoïque), Université des Sciences et Technologies de Lille, Villeneuve d'Ascq Cedex, F-59655, France.

Collaborators. R. Albani, A. Di Milia, K. Dorning, O. Fatka, T. Heuse, A. Le Hérissé, J. Li, M. Masiak, S.G. Molyneux, G. Playford, E. Raevskaya, C. Ribecai, C.V. Rubinstein, M. Stempien-Salek, L. Stricanne, M. Vanguetaine, M. Vecoli, A. Wallin, R. Wicander.

A major objective of the IGCP project 410 "The Great Ordovician Biodiversification Event" (1997-2002) was the assessment of biodiversity patterns of all taxonomic groups. In order to understand if the marine phytoplankton production in the Ordovician can be correlated to the invertebrate diversity, all data from Ordovician acritarchs have been compiled: some 700 articles have been published, describing over 250 genera (Servais, 1998) and over 1300 species and infraspecific taxa from Ordovician localities (Servais et al., 2004). The database of species described from the Ordovician Series includes over 1300 species of which most are established from localities of Gondwana and peri-Gondwana (some 400 species) and of Baltica (some 300 species), because most localities investigated are located in Europe and northern Africa. Over 250 new species have been described from the former Soviet Union, and nearly 100 from China. Data from North America and Australia add information to the global picture, while many other areas remain fairly or totally unstudied. In addition, most data on Ordovician acritarchs come from the Early to Middle Ordovician, publications of the Upper Ordovician are less abundant. Many taxonomic problems exist and the systematic position, as well as the spatial and temporal distribution of most species is not known. The elements to understand the diversity evolution of the Ordovician phytoplankton at a global scale are thus far from comprehensive. Global extinction events on the acritarch assemblages are not observed during the Early and Middle Ordovician, but at the end of the Upper Ordovician local changes have been reported: a great number of taxa disappeared before a new generation of taxa radiated during the beginning of the Silurian. More

precise information are provided at regional scales, especially from South China (Li in Servais et al., 2004) and North Africa (Vecoli in Servais et al., 2004).

Talk

Proposal of a new IGCP project: Ordovician Palaeogeography and Palaeoclimate : IGCP project no 503

T. Servais¹, D.A.T. Harper², J. Li³, A. Munnecke⁴, A.W. Owen⁵ & P.M Sheehan⁶

1, Paléontologie et Paléogéographie du Paléozoïque, UMR 8014 du CNRS, Université des Sciences et Technologies de Lille, Bâtiment SN5, 59655 Villeneuve d'Ascq, France. 2, Geologisk Museum, Øster Voldgade 5-7, 1350 København, Denmark. 3, Institute of Geology and Palaeontology, Chinese Academy of Sciences (NIGPAS), Chi-Ming-Ssu, Nanjing 210008, China. 4, Institut für Paläontologie, Universität Erlangen, Loewenichstraße 28, 91054 Erlangen, Germany. 5, Division of Earth Sciences, Centre for Geosciences, University of Glasgow, Gregory Building, Lilybank Gardens, Glasgow G12 8QQ, Scotland, UK. 6, Geology Department, Milwaukee Public Museum, 800 West Wells Street, Milwaukee, WI 53233, USA.

In October 2003, we proposed a new project, entitled "Ordovician Palaeogeography and Palaeoclimate: The impact of the changing palaeogeography and palaeoclimate on the major biotic changes through the Ordovician (Ordovician biodiversification, end-Ordovician extinction, Silurian radiation)" to the board of the International Geological Correlation Programme (IGCP). If accepted, our new project (n° 503) should run over five years from 2004 to 2008.

The project can be outlined as follows: arguably the most sustained rise in marine biodiversity took place during the Ordovician and the second largest mass extinction event took place close to the end of that Period, coincident with an episode of major climate fluctuation. The results of the very successful IGCP project n° 410 "The Great Ordovician Biodiversification Event" not only included the development of an improved globally- integrated biozonation for graptolites, conodonts and chitinozoans, but also generated biodiversity curves that have been constructed for all Ordovician fossil groups. Numerous questions arise from these results, for example how has changing palaeogeography affected the biodiversifications observed during the Ordovician, the extinctions at the end of the period, and the subsequent radiation in the Silurian. What was the influence of the climate on the major biotic changes through the Ordovician and Silurian? Following the work of the numerous regional teams and of the clade teams, that were established for each fossil group in IGCP project n° 410, we propose a new successor project in order to attempt to answer some of these questions (and generate others), and develop a better understanding of the environmental changes, e.g., global sea-level, temperature, oceanic circulation, climatic thresholds, etc. during the Early Palaeozoic. The new project will be developed in collaboration with the Subcommittee on Ordovician Stratigraphy (SOS) and with the Subcommittee on Silurian Stratigraphy (SSS).

The following working plan has been proposed: (1) the first year of the project will culminate in an assessment of all available information on ocean and climate modelling, as well as from the development of stable C- and O-isotopes in the Lower Palaeozoic; (2) the second year will focus on the evolutionary palaeoecology of the Early Palaeozoic, in order to understand the architecture of the ecosystems that developed and changed during the Ordovician biodiversification and extinction, and during the subsequent Silurian radiation and extinctions; (3) in the third year,

relationships will be sought between the changing palaeogeographical patterns and the changing biodiversities observed during the investigated intervals; (4) in the fourth year, we plan to compile all information on Early Palaeozoic events and stratigraphy, in order to improve the international standard time scale; (5) in the last year of our project, all information collected from the different regional teams should allow us to reconstruct Early Palaeozoic sea-level changes. At the end of our project, we also plan to produce a final synthesis bringing together the varied facets of the project and how they have addressed the objectives.

Five major meetings are scheduled between 2004 and 2008. Each of these meetings comprises indoor sessions, including the presentation of new results, and field excursions:

In 2004, if accepted, the official opening meeting of the new IGCP project will be organised at the Universität Erlangen, Germany, September, 1-3, 2004, just following the International Geological Congress at Florence, Italy. This meeting will be focused on Early Palaeozoic Climate Modelling and isotope geochemistry. The associated geological excursion will cover the Ordovician of Öland and the Silurian of Gotland (Sweden).

The second major meeting will be held in 2005 at the Milwaukee Public Museum (Wisconsin, USA) and focus on the ecological evolution in the Early Palaeozoic. The geological excursions will be organized in the Ordovician of the American mid-Continent (Kentucky, Ohio, etc.). We plan to organize this conference just before or after the North American Paleontological Convention meeting.

In 2006, a major meeting of the new IGCP project will be held at Glasgow University, Scotland, UK, and concentrate on palaeo(bio)geography, and the relation between Early Palaeozoic biodiversity and climatic belts. A geological excursion to the Lower Palaeozoic of Scotland and northern England will include a visit to the GSSP of the base of the Silurian at Dob's Linn and to Scottish Silurian inliers in Southern Uplands and Pentland Hills.

The Nanjing Institute of Geology and Palaeontology, Academia Sinica, China, will organize the 10th ISOS in 2007. Jointly with the Ordovician Meeting, the Silurian Subcommittee will also hold its International Symposium. An IGCP related symposium on Early Palaeozoic events is scheduled at Nanjing, as well as several geological excursions to Lower Palaeozoic sections of southern China, to Ordovician to Early Silurian sections of the Upper Yangtze Platform.

The closing meeting of our project should take place in 2008 at the Université des Sciences et Technologies de Lille, France. This meeting is dedicated to the reconstruction of Early Palaeozoic sea-level changes. Geological excursions to Lower Palaeozoic sections of France and Belgium will be organised.

A series of additional meetings are already scheduled (Argentina, Spain, Czech Republic, Estonia, etc.).

Talk

Similar patterns of vertical distribution in Silurian acritarchs: another tool for the reconstruction of palaeoenvironmental changes in the Ludlow of Gotland, Sweden

Ludovic Stricanne

Paléontologie et Paléogéographie du Paléozoïque, UMR 8014 du CNRS, Université des Sciences et Technologies de Lille, Bâtiment SN5, 59655 Villeneuve d'Ascq, France.
ludovic.stricanne@ed.univ-lille1.fr

Most of the investigation on Palaeozoic acritarchs concern the biostratigraphy although palaeoecological approaches still need to be developed. In the Ludlow (Late Silurian) of Gotland (Baltic Sea), such an approach has been tentatively introduced on two abundant and diversified acritarch genera, *Michrhystridium* Deflandre 1937 and *Veryhachium* Deunff 1954. The vertical distribution of these genera have been quantitatively investigated and show large changes in their development and of their species. In the Gorstian, most of the taxa are well ornamented (long processes) and the morphological diversity is high, whereas upper Ludfordian assemblages are extremely reduced in diversity, with morphotypes of short process length. At the Early and Late Ludfordian boundary, both genera show a continuous shift to more simple morphologies, associated with palaeoenvironmental and palaeoclimatic changes (arid conditions) as deduced from the facies occurrences and a large $\delta^{13}\text{C}$ isotope excursion, respectively. At the end of the Ludfordian, the throwback to normal marine conditions provides morphotypes with an enhanced ornamentation. Similar trends in the frequency of morphotypes are also observed in a Gorstian lateral-transect. Thus, these results allow some considerations regarding the palaeoecological properties of some acritarch taxa, frequently used for biostratigraphical purpose. The quantitative analysis of large populations may provide an accurate tool for the potential use of selected acritarch taxa for the reconstructions of Silurian palaeoenvironments.

Talk (cancelled)

Sea levels, paleoclimate and hydrocarbon systems in Bolivia

Ian Troth

Southampton Oceanography Centre, European Way, Southampton, SO14 3ZH, England.
it598@soc.soton.uk

In Bolivia there is a world class reservoir (Huamampampa Sandstone) and source rock (Los Monos Shales). Producing fields are in the sub-Andean zone where they occur in a structurally complex thrust terrain. Seismic recognition in the subsurface of the reservoir is difficult because of this structural complexity. However, success has been achieved using palynology. A bloom of the acritarch (marine phytoplankton) *Evittia sommerii* occurs in the lowest Los Monos Shales, immediately above the Huamampampa Sandstone. The bloom only occurs at this stratigraphic level and represents an influx of this distinctive acritarch into the basin. This event has not been studied before at surface outcrop. Surface exposures of the Huamampampa-Los Monos interval have been investigated over two field seasons and the rock samples collected have revealed the precise stratigraphic extent of the bloom. This data will be used for bio-steering purposes during drilling in Bolivia. In Devonian times, Bolivia was in the high latitude, coldwater Malvinokaffric province. The *Evittia* bloom is monospecific and is coincident with a peak of prasinophyte algae (leiospheres) that are generally associated with deeper water. The presence of these

two palynomorphs suggests there was a sea level rise possibly associated with an influx of warm water from lower latitudes. An influx of warm water would have contributed to the breakdown of the Malvinokaffric realm in the Mid-Devonian times.

Talk

Biodiversity patterns of Ordovician marine microphytoplankton: biological and palaeoecological meaning

Marco Vecoli

Laboratoire de Paléontologie, Université de Bretagne Occidentale and CNRS. Street. 6, Avenue Le Gorgeu, P.O. BOX 809, Brest 29285, France. marco.vecoli@univ-brest.fr; marco_vecoli@hotmail.com

Calculation of evolutionary rates and diversity curve at specific level, form the basis for discussing the influence of global palaeoenvironmental perturbations on the evolution of organic-walled microphytoplankton in northern Gondwana during latest Cambrian through Ordovician times.

The most important Cambro-Ordovician acritarch bio-events are short periods of diversification, which also correspond to introduction of morphological innovations, observed in latest Cambrian and earliest Tremadoc, late Tremadoc, early Arenig, basal Llanvirn, and latest Ashgill, and an important extinction phase in the early Caradoc. Overall, acritarch diversity increased from the basal Ordovician up to the middle Llanvirn, then declined in the early and middle Caradoc. During Ashgill times, the assemblages are poorly diversified at the generic level as a result of a combined effect of sea level drawdown and onset of glacial conditions, but no major extinction event is observed in connection with the end-Ordovician biotic crisis. The peak in acritarch diversity during Middle Ordovician times appears to be correlated to maximum spread of palaeogeographical assembly. Acritarch dynamics appear largely uncorrelated to second order sea-level oscillations; the primary abiotic controls on acritarch evolution were palaeogeographical and the associated palaeoceanographic changes (especially during Middle Ordovician), and the end-Ordovician palaeoclimatic shift.

The acritarch fossil record provides important information on the evolution of oceanic primary producers, however, the relationships between acritarch diversity, oceanic productivity, and evolution of invertebrate animals are proving much more complex than previously thought. In particular, the hypothesis of a causal relationship between changes in acritarch diversity and metazoan evolution in the Palaeozoic is not supported by our data.

Talk

An overview of Devonian acritarchs and prasinophytes

Reed Wicander

Department of Geology, Central Michigan University, Mount Pleasant, Michigan U.S.A.
48859. reed.wicander@cmich.edu

Although there appears to be a fairly extensive literature on Devonian acritarchs and prasinophytes during the past thirty years, their study has, in fact, been somewhat uneven, both stratigraphically and geographically. There is, however, a growing body of literature in which well-preserved acritarch and prasinophyte assemblages from independently dated surface and subsurface sections are described allowing for regional and global zonations and correlations.

Most Devonian published acritarch and prasinophyte studies have been concentrated primarily in North America and Europe, followed by South America, Australia, and Asia. It is becoming increasingly clear that Lower, Middle, and Upper Devonian acritarch and prasinophyte assemblages are fairly distinct from each other. The literature indicates the least is known for the Lower Devonian both in diversity of the acritarch and prasinophyte assemblages and geographic distribution, followed by greater coverage for the Middle Devonian, with the greatest attention focused on the Upper Devonian, particularly North America and Europe.

The distribution of Devonian acritarchs and prasinophytes is presumably influenced by the same factors affecting modern phytoplankton taxa, namely light, temperature, salinity, nutrients, water depth, and surface currents. There appears to be a correlation between increasing diversity and change in morphotypes with increasing distance from the paleoshoreline as well as a change in the spore/acritarch ratio that is related to paleoshoreline proximity.

Before any meaningful conclusions can be made concerning the effect of fluctuating acritarch and prasinophyte diversity patterns on the Devonian marine ecosystem and climate, the role played by ecology, biogeography, taxonomy, and taphonomy must be addressed and analyzed.

Talk