

The Moment Generating Function of the Binomial Distribution

Consider the binomial function

$$(1) \quad b(x; n, p) = \frac{n!}{x!(n-x)!} p^x q^{n-x} \quad \text{with} \quad q = 1 - p.$$

Then the moment generating function is given by

$$(2) \quad \begin{aligned} M_x(t) &= \sum_{x=0}^n e^{xt} \frac{n!}{x!(n-x)!} p^x q^{n-x} \\ &= \sum_{x=0}^n (pe^t)^x \frac{n!}{x!(n-x)!} q^{n-x} \\ &= (q + pe^t)^n, \end{aligned}$$

where the final equality is understood by recognising that it represents the expansion of binomial. If we differentiate the moment generating function with respect to t using the function-of-a-function rule, then we get

$$(3) \quad \begin{aligned} \frac{dM_x(t)}{dt} &= n(q + pe^t)^{n-1} pe^t \\ &= npe^t (q + pe^t)^{n-1}. \end{aligned}$$

Evaluating this at $t = 0$ gives

$$(4) \quad E(x) = np(q + p)^{n-1} = np.$$

Notice that this result is already familiar and that we have obtained it previously by somewhat simpler means.

To find the second moment, we use the product rule

$$(5) \quad \frac{d^2 M_x(t)}{dt^2} = u \frac{dv}{dx} + v \frac{du}{dx}$$

to get

$$(6) \quad \begin{aligned} \frac{d^2 M_x(t)}{dt^2} &= npe^t \{(n-1)(q + pe^t)^{n-2} pe^t\} + (q + pe^t)^{n-1} \{npe^t\} \\ &= npe^t (q + pe^t)^{n-2} \{(n-1)pe^t + (q + pe^t)\} \\ &= npe^t (q + pe^t)^{n-2} \{q + npe^t\}. \end{aligned}$$

Evaluating this at $t = 0$ gives

$$(7) \quad \begin{aligned} E(x^2) &= np(q+p)^{n-2}(q+np) \\ &= np(q+np). \end{aligned}$$

From this, we see that

$$(8) \quad \begin{aligned} V(x) &= E(x^2) - \{E(x)\}^2 \\ &= np(q+np) - n^2p^2 \\ &= npq. \end{aligned}$$

Theorems Concerning Moment Generating Functions

In finding the variance of the binomial distribution, we have pursued a method which is more laborious than it need be. The following theorem shows how to generate the moments about an arbitrary datum which we may take to be the mean of the distribution.

$$(9) \quad \text{The function which generates moments about the mean of a random variable is given by } M_{x-\mu}(t) = \exp\{-\mu t\}M_x(t) \text{ where } M_x(t) \text{ is the function which generates moments about the origin.}$$

This result is understood by considering the following identity:

$$(10) \quad M_{x-\mu}(t) = E\{\exp\{(x-\mu)t\}\} = e^{-\mu t}E(e^{xt}) = \exp\{-\mu t\}M_x(t).$$

For an example, consider once more the binomial function. The moment generating function about the mean is then

$$(11) \quad \begin{aligned} M_{x-\mu}(t) &= e^{-npt}(q+pe^t)^n \\ &= (qe^{-pt} + pe^te^{-pt})^n \\ &= (qe^{-pt} + pe^{qt})^n. \end{aligned}$$

Differentiating this once gives

$$(12) \quad \frac{dM_{x-\mu}(t)}{dt} = n(qe^{-pt} + pe^{qt})^{n-1}(-pqe^{-pt} + qpe^{qt}).$$

At $t = 0$, this has the value of zero, as it should. Differentiating a second time according to the product rule gives

$$(13) \quad \frac{d^2M_{x-\mu}(t)}{dt^2} = u(p^2qe^{-pt} + q^2pe^{qt}) + v\frac{du}{dt},$$

where

$$(14) \quad \begin{aligned} u(t) &= n(qe^{-pt} + pe^{qt}) \quad \text{and} \\ v(t) &= (-pqe^{-pt} + qpe^{qt}). \end{aligned}$$

At $t = 0$ these become $u(0) = n$ and $v(0) = 0$. It follows that

$$(15) \quad V(x) = n(p^2q + q^2p) = npq(p + q) = npq,$$

as we know from a previous derivation.

Another important theorem concerns the moment generating function of a sum of independent random variables:

$$(16) \quad \text{If } x \sim f(x) \text{ and } y \sim f(y) \text{ be two independently distributed random variables with moment generating functions } M_x(t) \text{ and } M_y(t), \text{ then their sum } z = x + y \text{ has the moment generating function } M_z(t) = M_x(t)M_y(t).$$

This result is a consequence of the fact that the independence of x and y implies that their joint probability density function is the product of their individual marginal probability density functions: $f(x, y) = f(x)f(y)$. From this, it follows that

$$(17) \quad \begin{aligned} M_{x+y}(t) &= \int_x \int_y e^{(x+y)t} f(x, y) dy dx \\ &= \int_x e^{xt} f(x) dx \int_y e^{yt} f(y) dy \\ &= M_x(t)M_y(t). \end{aligned}$$